

Contents lists available at ScienceDirect

# Marine Pollution Bulletin



journal homepage: www.elsevier.com/locate/marpolbul

# Ecological status and sources of anthropogenic contaminants in mangroves of the Wouri River Estuary (Cameroon)



Marco Fusi<sup>a,\*</sup>, Gian Maria Beone<sup>b</sup>, Nicoleta Alina Suciu<sup>b</sup>, Angela Sacchi<sup>b</sup>, Marco Trevisan<sup>b</sup>, Ettore Capri<sup>b</sup>, Daniele Daffonchio<sup>a</sup>, Ndongo Din<sup>c</sup>, Farid Dahdouh-Guebas<sup>d,e,1</sup>, Stefano Cannicci<sup>f,g,1</sup>

<sup>a</sup> King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences & Engineering Division, Thuwal 23955-6900, Saudi Arabia

<sup>b</sup> Institute of Agricultural and Environmental Chemistry, Università Cattolica del Sacro Cuore, Via Emilia Parmense 84, 29122 Piacenza, Italy

<sup>c</sup> The University of Douala, Faculty of Science, Department of Botany, P.O. Box 8948, Douala, Cameroon

<sup>d</sup> Laboratory of Systems Ecology and Resource Management, Department of Organism Biology, Faculty of Sciences, Université Libre de Bruxelles – ULB, Avenue F.D. Roosevelt 50, CPI 264/1, B-1050 Brussels, Belgium

<sup>e</sup> Laboratory of Plant Biology and Nature Management, Department of Biology, Faculty of Sciences and Bio-engineering Sciences, Vrije Universiteit Brussel – VUB, Pleinlaan 2, B-1050 Brussels, Belgium

<sup>f</sup> The Swire Institute of Marine Science and The School of Biological Sciences, The University of Hong Kong, Hong Kong

<sup>g</sup> Department of Biology, University of Florence, Via Madonna del Piano 6, Sesto Fiorentino, Italy

#### ARTICLE INFO

Article history: Received 11 December 2015 Received in revised form 23 June 2016 Accepted 28 June 2016 Available online 6 July 2016

Keywords: Mangrove Contaminants Macrobenthos Cameroon Wouri Estuary Sediment

# 1. Introduction

# ABSTRACT

Mangroves are critically threatened by human activities, despite the important ecosystem functions and services they provide. Mangroves in Cameroon represent no exception to the worldwide trend of mangrove destruction, especially around Douala, on the Wouri river estuary. In two sites around Douala, we assessed the presence of sterols, PAHs, PCBs, DEHP, DDT and its metabolite p,p'-DDE and potentially toxic metals in sediment samples. As a proxy of ecological quality, we measured the diversity and abundance of macrobenthos assemblages. We detected p,p'-DDE contamination, with concentrations higher than 3  $\mu$ g kg<sup>-1</sup> in 16 out of 26 samples which were attributed to recent widespread use of DDT. The detection of sterols revealed faecal contamination. Significant sensitivity of the macrobenthos to contaminants was revealed, with possible implications on the overall mangrove vulnerability to climate change and on the provision of ecosystem services to local populations.

© 2016 Elsevier Ltd. All rights reserved.

Mangroves in Africa cover over 3.2 million ha, corresponding to about 20% of their global coastline coverage, with approximately 1.5 million ha located along the Atlantic coast (Giri et al., 2011; Massó i Alemán et al., 2010; Spalding et al., 2010; UNEP, 2007). As a consequence of enormous anthropogenic pressure and multiple threats, western African mangroves have declined by >25% over the past 25 years (Friess and Webb, 2014; Giri et al., 2011). Cameroon harbours approximately 2000 km<sup>2</sup> of mangroves, distributed along the coast of the Guinean gulf (Giri et al., 2011). Although mangroves contribute considerably to the social and economic well-being of the Cameroonian coastal inhabitants, their total surface area has decreased by 30% in 20 years (Spalding et al., 2010), mainly due to rapid and uncontrolled urbanization around Douala (Din et al., 2002; Ellison and Zouh, 2012; Nfotabong-Atheull et al., 2013). With a population of >2 million people, Douala is the largest

city in Cameroon and exerts a huge pressure on the nearby mangroves, with uncontrolled sewage discharge detrimentally affecting the whole ecosystem (Simon and Raffaelli, 2012).

Douala is also one of the major shipping ports in the Guinea Gulf that serves the entire central Africa and refuels oil tankers to export locally extracted oil, another significant anthropogenic impact on the Wouri River estuary mangroves (Alemagi, 2007; Duke, 2016; Price et al., 2000; Van De Walle, 1989). Due to the lack of policy regulation in the management of Cameroonian coastal ecosystems, sand mining and wood harvesting also play an important role in reducing mangrove biodiversity and provision of ecosystem services (Ellison and Zouh, 2012; Nfotabong-Atheull et al., 2011).

Although these multiple impacts threaten Wouri River estuary forests, the major socio-economic activity associated with mangroves for local people is in fact still artisanal fishing, with landings estimated between 76 and 106 tons per year (Gabche, 1997). Fisheries play a significant role in small-scale commercial activities and they are vital in providing a source of protein and income for coastal communities (Nfotabong-Atheull et al., 2009). Thus, the modification of both abundance and diversity of mangrove species and the deterioration of

<sup>\*</sup> Corresponding author.

E-mail address: marco.fusi@kaust.edu.sa (M. Fusi).

<sup>&</sup>lt;sup>1</sup> These authors equally contributed to the paper.

water quality, due to urban and industrial activities, will surely have detrimental consequences on the well-being of local communities (Alemagi, 2007; Nfotabong-Atheull et al., 2009, 2011). Last but not least, vulnerability to climate change, and especially to sea level rise, proved to be exacerbated by the high level of anthropogenic pressure on the Wouri River estuary mangroves (Ellison and Zouh, 2012). In particular, purported impacts are (i) the increase of frequency and duration of the tidal inundation that may cause the death of the mangrove trees (exceeding the species-specific physiological thresholds; Ball, 1988), (ii) the impact on the inland fresh groundwater with saline intrusion and contaminants dispersal in the intertidal systems (Woodroffe et al., 2016), and (iii) the change of the topography and hydrology of the sediment (Lovelock et al., 2015).

Our aim was to perform the first baseline study on the ecological status and pollution of the strongly impacted Wouri River estuary mangroves, collecting data on both the presence of anthropogenic pollutants in sediments and the structure and diversity of macrobenthic populations as a proxy for healthy ecosystem functioning (Cannicci et al., 2008, 2009). To assess the level of chemical pollution, we targeted the major anthropogenic compounds usually found in peri-urban impacted mangrove forests world-wide, i.e. organochlorine compounds, such as DDT and its metabolites, phthalates such as bis(2ethylhexyl)phthalate (DEHP), polycyclic aromatic hydrocarbons (PAHs), polychlorobyphenyls (PCBs), heavy metals and sterols (Bayen et al., 2005, 2016; Lewis et al., 2011; MacFarlane et al., 2007; Peters et al., 1997; Vane et al., 2009). Compounds such as coprostanol ( $5\beta$ (H)-Colestan-3 β-ol) can be used in conjunction with other sterols to determine the relative abundance of sewage in sediments. Coprostanol, in particular, is a faecal sterol generated by microbial activity on cholesterol and is considered as a chemical marker of faecal contamination, especially from humans (Bull et al., 2002; Fattore et al., 1996; Mudge et al., 1999; Peng et al., 2005; Sherwin et al., 1993).

Diversity and abundance of crab and mollusc populations were recently shown to be key-determinants of the maintenance of mangrove ecosystem function and services (Cannicci et al., 2008; Duke et al., 2007; Lee, 2008), such as the provision of nursery sites for fish stocks which is of great importance for the local economy. Crabs and molluscs form an important link between primary detritus at the base of the food web and consumers at higher trophic levels (Sousa and Dangremond, 2011). By consuming litter, crabs can promote nutrient mineralization and recycling within the forest. Furthermore, their bioturbation activities undoubtedly alter the physico-chemical characteristics of soil (Kristensen, 2008) and enhance below-ground organic carbon retention (Andreetta et al., 2014). Finally, since mangrove macrobenthos diversity and functioning are known to be strongly impacted by contaminants (see Cannicci et al., 2009; Bartolini et al., 2011; Penha-Lopes et al., 2011 for east African mangrove benthos), their abundance and diversity is useful in assessing the degree of bioavailability of anthropogenic pollutants and the actual impact on the biological components.

#### 2. Material and methods

# 2.1. Area description

The study was carried out in two peri-urban mangrove forests located at different sites along the Wouri estuary: Wouri Bridge forest (4°4′ 19.10880″N; 9°42′5.81312″ E, hereafter WB) and Bois des Singes forest (4°0′49.67706″N; 9°40′ 28.10325″E, hereafter BS), located north-west and south-east of Douala, respectively (Fig. 1), with a distance between them of 11 km. Both of these stands are at about 10 ha in extension and are largely affected by the uncontrolled expansion of urban areas due to the rapidly increasing population of Douala city (Simon and Raffaelli, 2012). Thus, they are representative sites to assess the possible presence of pollutants in peri-urban mangroves. The climate of the region belongs to the Equatorial regime (Din and Baltzer, 2008), characterized by a long rainy season (March–November) and a short dry season (December–February). Heavy rainfall (approximately 4000 mm per year), stable high temperatures (annual average temperature is 26.7 °C) and high humidity throughout the year (approaching 100%) are typical for this region. The tidal regime is semi-diurnal with an average amplitude of 2.5 m. Soils are grey or black muds, of silty, sandy or clayey texture, derived from fluvial sediments relatively rich in organic matter with a high C:N ratio due to the reduced biological activity (Campo and Darius, 2004). Annual salinity variation in the region ranges between 0 and 20‰. During the long monsoon season, mangrove water salinity is consistently <10‰. During the dry season, salinity varies between 4 and 20‰ (Din and Baltzer, 2008).

According to a survey by Saenger and Bellan (1995), the floristic composition of Wouri Bridge forest, a 40 year old stand, is dominated by *Avicennia germinans*, *Rhizophora racemosa*, *Rhizophora mangle* and *Rhizophora harrisonii* and the mangrove associate *Pandanus* sp. Bois des Singes is an older stand and has a different floristic composition, represented only by the three *Rhizophora* species listed above. Hereafter, the belts studied will be referred as Avicennia belt, Pandanus belt and Rhizophora belt.

In these systems the faunal composition includes vertebrates, such as mammals like the sirenian species *Trichechus senegalensis* and several cetaceans. There are birds, of which particular significance is the high abundance of african skimmers, grey pranticoles, open-billed storks and common green shanks, reptiles such as four species of sea turtles (see Diop et al., 2014), and fishes (many of them of commercial importance), and a wide range of invertebrates, mainly crabs (belonging to the families Sesarmidae and Ocypodidae) and molluscs (extensively described by Ngo-Massou et al., 2012), which constitute the bulk of ben-thic diversity in the these mangrove ecosystem.

#### 2.2. Sediment sampling

A total of 20 sediment samples were collected in September 2009 in the two mangrove systems for trace metals and organic compounds analysis, choosing five random replicates in each forest. Five samples from BS and five from WB were taken from the upper layer of superficial sediment (0–10 cm) and other five samples from BS and five from WB were taken from the layer underneath (11–20 cm). The sediment cores were collected using an Eijkelkamp Multisampler<sup>TM</sup> piston corer (10 cm diameter). The samples were then placed in glass jars, covered with aluminium foil and immediately transferred to a portable freezer and stored at -20 °C until analysis.

#### 2.3. Analytical methods

## 2.3.1. Solvents, chemicals and standards

The solvents used were acetone, hexane, dichloromethane and isooctane, obtained from Sigma Aldrich and Fluka Co., Steinheim, Germany. Standard reference materials for trace metals analysis were supplied by the Community Bureau of Reference Sample (BCR): Certified Reference Materials CRM 277 and CRM 320 and 142 R. Analytical standards for a mixture of PCBs (IUPAC nr. 28, 52, 101, 118, 138, 153, 180), a mixture of PAHs (anthracene, benzo[*a*] anthracene, benzo [jbk]fluoranthene, benzo [a] pyrene, benzo [ghi] perylene, chrysene, fluoranthene, indeno[1,2,3-cd]pyrene, phenanthrene, pyrene), bis(2ethylhexyl)phthalate (DEHP) and the internal standards anthracened<sub>10</sub> and perylene-d<sub>12</sub> were purchased from Dr. Ehrenstorfer GmbH, Augsburg, Germany. Analytical sterol standards, Coprostan-3-ol, 5 α-Cholestan-3 $\beta$ -ol, cholesterol and 5  $\beta$ -Cholestan-3 $\alpha$ -ol, analytical Bis(trimethylsilyl)trifluoracetamide (BSTFA) with 1% trimethylchlorosilane (TMCS), used for sterols derivatisation, and analytical standards for 1,1,1-trichloro-2,2-di(4-chlorophenyl)ethane (DDT) and 1,1-bis-(4-chlorophenyl)-2,2-dichloroethene (p-p'DDE) were purchased from Sigma Aldrich and Fluka Co, Steinheim, Germany.



Fig. 1. Study sites. A) Continental overview, B) Woury Estuary where Douala is located, C) Wouri Bridge mangrove forest (WB) and D) Bois de singes mangrove forest (BS). Black squares indicate the exact location where the study was carried on (Image source: Google Earth).

#### 2.3.2. Trace metals analysis

All analyses were performed one month after sampling. Trace element concentrations of Mn, Pb, Cr, Cu, Zn, V and Ni were determined by inductively coupled plasma optical emission spectrometry (ICP-OES. Perkin Elmer Optima 2100 DV spectrometry, Massachusetts, USA). Concentrations of Co. As. Se. Mo. Cd. Sn. Sb and Tl were determined by inductively coupled plasma mass spectrometry (ICP-MS, Agilent Technologies mod. 7700× with Octopole Reaction System ORS, Santa Clara, USA) following aqua regia digestion according to Bettinelli et al. (2000). Mercury was determined by an automatic solid Hg analyser AMA 254. Certified and experimental values exhibited consistent values, recoveries ranging between 93 and 106% with repeatability better than 8% using CRM 277 'Estuarine Sediment', CRM 320 'River Sediment' and CRM 142 'Light Sandy Soil'. Certified soils and sediments were supplied by The Community Bureau of Reference Sample (BCR -IRMM, Joint Research Center, Retieseweg, B-2440, Geel, Belgium). CMR samples were handled according to the supplier's specifications. The MDL value was also calculated by analysing the blanks prepared on different working days. This parameter was assumed to be three times the blanks' standard deviation (See Table S1 in the supplementary material).

## 2.3.3. Organic compounds analysis

For the extraction of PAHs (anthracene, benzo[*a*] anthracene, benzo [*jbk*]fluoranthene, benzo [*a*]) pyrene, benzo [ghi] perylene, chrysene, fluoranthene, indeno[1,2,3- cd]pyrene, phenanthrene, pyrene, PCBs (IUPAC nr. 28, 52, 101, 118, 138, 153, 180), DEHP, DDT and its metabolite, samples were treated according to Zaccone et al. (2009). After extraction with Soxhlet using a hexane (80%) and acetone (20%) mixture and concentration of the extracts using a Buchi B-811 Rotavapor, the obtained solutions were divided into two equal parts. A 5 ml aliquot of the extract was evaporated under a gentle flow of nitrogen, recovered with 0.5 mL of hexane containing the internal standards anthracene d<sub>10</sub> (1.14 mg l<sup>-1</sup>) and perylene d<sub>12</sub> (1.05 mg l<sup>-1</sup>), centrifuged and analysed with GC–MS to determine the presence of benzo[*jbk*]fluoranthene (*m*/*z* 252), benzo[*a*]pyrene (*m*/*z* 252), benzo [*ghi*]perylene (*m*/*z* 276), and indeno[1,2,3- cd]pyrene (*m*/*z* 276), DEHP (*m*/*z* 149), DDT (*m*/*z* 235) and p,p'-DDE (*m*/*z* 246). The remaining 5 mL of the extracts were cleaned-up through a Florisil column, evaporated under a gentle flow of nitrogen, dissolved in 0.5 mL of hexane containing the internal standards anthracene d<sub>10</sub> (1.14 mg l<sup>-1</sup>) and perylene d<sub>12</sub> (1.05 mg l<sup>-1</sup>), and then analysed to determine the presence of anthracene (*m*/*z* 178), chrysene (*m*/*z* 228), benzo [*a*]anthracene (*m*/*z* 228), phenanthrene (*m*/*z* 178), pyrene (*m*/*z* 292 for nr. 52, *m*/*z* 326 for nr. 101and nr. 118, *m*/*z* 360 for nr.138 and nr. 153 and *m*/*z* 394 for nr.180).

GC–MS analysis was performed according to Zaccone et al. (2009). Total Ion Monitoring (TIM) and Selected Ion Monitoring mode (SIM) were used for identification and quantification of substances. Overall concentration of PAHs and PCBs in sediment is the sum of the 12 PAHs and 7 PCBs, respectively, analysed individually (mean of five samples for each site) expressed on a dry weight basis.

The sterols were analysed following the method proposed by Froehner et al. (2009), which includes Soxhlet extraction, clean-up with a silica-aluminium column, derivatisation and detection and quantification by GC–MS. The LODs were 0.2  $\mu$ g kg<sup>-1</sup> for coprostan-3-ol and 0.4  $\mu$ g kg<sup>-1</sup> for 5  $\alpha$ -Cholestan-3 $\beta$ -ol, Cholesterol and 5  $\beta$ -Cholestan-3 $\alpha$ -ol. The recoveries of sterols/stanols were between 65 and 80%.

The sediment total organic carbon (TOC) and total nitrogen (TN) in pore water were determined using the standard methods recommended by SSSA (Sparks et al., 1996).

# 2.3.4. Quality control

Calibration curves, prepared by dilution of stock solution with hexane, for PAHs and PCBs and DEHP were obtained at concentrations between 0.01 and 0.2 mg  $l^{-1}$  and 0.1 and 2 mg  $l^{-1}$ , respectively, using anthracene  $d_{10}$  (1.14 mg l<sup>-1</sup>) and pervlene d12 (1.05 mg l<sup>-1</sup>) as internal standards. The calibration curves for p,p'-DDE and DDT were created at concentrations between 0.01 and 1 mg  $l^{-1}$ , using anthracene d10  $(1.14 \text{ mg } l^{-1})$  and pervlene d12  $(1.05 \text{ mg } l^{-1})$  as internal standards. Calibration curves for sterols were obtained at concentrations between 0.05 and 5 mg  $l^{-1}$  by the dilution of stock solution with isooctane. Recovery experiments ( in triplicate) for PAH and PCBs were performed on a Certificate Reference Material (CRM) IMEP-21 obtained from the European Commission-JRC-IRMM with obtained values between 84 and 130% for PAHs and between 70 and 115% for PCBs and standard deviation, in all the cases, <20%. The limit of detection (LOD) was 2  $\mu g \, kg^{-1}$ for PAHs and 0.5  $\mu$ g kg<sup>-1</sup> for PCBs. Due to the fact that DEHP was present in the extraction solvents and other materials used for extraction, three blank extractions were undertaken during the DEHP extraction process for all the samples. The DEHP sample concentrations were then corrected based on the daily blank extraction values. A Recovery experiment (in triplicate) for DEHP was performed on a sewage sludge sample by sample contamination at a concentration of 1 mg kg<sup>-1</sup>, with obtained values of 90% and standard deviation of 5%. The LOD for DEHP was 5  $\mu$ g kg<sup>-1</sup> whereas for p,p'-DDE it was 0.5  $\mu$ g kg<sup>-1</sup>. The recoveries of sterols/stanols were between 65 and 80% with a standard deviation <5%, whereas the LODs were 0.2 µg kg<sup>-1</sup> for coprostan-3-ol and 0.4  $\mu$ g kg<sup>-1</sup> for 5  $\alpha$ -Cholestan-3 $\beta$ -ol, Cholesterol and 5  $\beta$ -Cholestan- $3\alpha$ -ol.

#### 2.4. Macrobenthos survey

In both forests, the surveys were carried out during three consecutive spring tides in September - October 2009. In each forest, two random transects (100-500 m apart) were established in each vegetation belt following a nested design. Along each transect, three  $2 \times 2$  m quadrats were randomly sampled to assess the abundance and density of the brachyuran and molluscan populations. The surveys were replicated for each spring tide period, at the same time of the day for the same time of observation in agreement with the methods described in Skov et al. (2002). Based on the complexity of the habitat and the diverse behaviour of the study species, different sampling techniques were used to assess the abundance of the various groups of macrofauna. Due to their high densities, molluscs were counted in a sub-quadrat of  $50 \times 50$  cm placed within the sampling quadrat. Small sesarmids were counted visually throughout the quadrats. Large sesarmids were assessed by counting the number of operational burrows within the quadrats, since previous studies in South Africa and Kenya have clearly shown that these refuges are occupied by single crabs (Berti et al., 2008; Emmerson, 2001; Fratini et al., 2000; Skov et al., 2002). In order to refine the evaluation of crab and mollusc numbers, and due to the accumulation of leaf litter obscuring crabs, after observation for 1 h in every quadrat we removed fallen leaves and logs to count the hidden specimens. Furthermore we measured temperature, pH and conductivity of the sediment water for each plot in each location using an Acorn pH 6 meter probe (Oakton Instruments).

#### 2.5. Statistical analyses

A non-metric multidimensional scaling ordination (nMDS) was performed on the basis of a Bray-Curtis dissimilarity matrix calculated on untransformed data to visualize patterns of macrobenthic composition across sites. Furthermore, a PERMANOVA (Anderson et al., 2008) was used to test the null hypothesis of no differences in macrobenthos assemblages and temperature, pH and conductivity across the factor Site (fixed, orthogonal, two levels: WB – Wouri Bridge and BS – Bois des Singes), Belt (fixed, orthogonal three levels: *Rhizophora, Pandanus* and *Avicennia*) and Transect (random, nested in Site, 2 levels). PERMANOVA was also used to test the null hypothesis that there were no differences in contaminants, with the factors Site (fixed, orthogonal, two levels: WB – Wouri Bridge and BS – Bois des Singes), Belt (fixed, orthogonal three levels: *Rhizophora, Panda-nus* and *Avicennia*) and Sampling Depth (fixed, orthogonal two levels: 0–10 and 11–20 cm). In the statistical analysis of contaminants, all values were normalised as performed by Dafforn et al. (2012), and Spearman's correlation was performed in order to eliminate covariate variables. For contaminants, the Euclidean distance was used to calculate the dissimilarity matrix. DistLM Analysis was performed to test the significant relationship between the ecological and anthropogenic factors, and the macrobenthos assemblages.

#### 3. Results

#### 3.1. Sediment analysis

Concentrations of ten PAHs (Fig. 2, supplementary table S2) and six PCBs congeners (Table 1) were lower than 300  $\mu$ g kg<sup>-1</sup> and 20  $\mu$ g kg<sup>-1</sup> of sediment dry weight respectively. No statistical differences were recorded among sites, belts and sampling depths. Similarly, metals (Fig. 2, supplementary table S3) and DEHP (Table 2) concentrations were consistently lower than 1500  $\mu$ g kg<sup>-1</sup>. DDT was absent, whereas its metabolite p-p'DDE (Table 2) was found at concentrations higher than  $3 \,\mu g \, kg^{-1}$  in 16 samples. Furthermore, no statistical differences among the samples or between sites, belts and depths were observed. Cholesterol and 5  $\alpha$ -cholestan-3 $\beta$ -ol concentrations were lower than 1000  $\mu$ g kg<sup>-1</sup> (Fig. 2, supplementary table S4), while concentrations of coprostan-3-ol and 5  $\beta$ -cholestan-3 $\alpha$ -ol were lower than 4000  $\mu g \ kg^{-1}$  (with the exception of two samples). No statistical differences were detected among sites, belts and sampling depths. Interestingly, Coprostanol and 5  $\beta$ -cholestan-3 $\alpha$ -ol were not found in the deepest layers of sediment of any of the belts examined at the Wouri Bridge site.

Ratios between different sterols are presented in Table 3 and they provide information concerning the source of contamination (according with Froehner et al., 2009), revealing widespread high disturbance in both forests.

#### 3.2. Macrobenthic assemblages

The two forests were similar in terms of temperature, sediment water pH and sediment water conductivity (PERMANOVA, n = 25, F = 0.42, df = 1,25, p > 0.05; Table 4), as well as total N and total OC (PERMANOVA, n = 28, F = 0.89, df = 2,28, p > 0.05; Table 4).

In accordance with the species listed by Ngo-Massou et al. (2012) and the reference list edited by Ng et al. (2008) and Manning and Holthuis (1981), we individuated seven species of sesarmid and two species of mollusc inhabiting the forest. Within Sesarmidae, with an average density of 1.5 individual per square meter, we identified Perisesarma kamermanni, Perisesarma huzardi, Metagrapsus curvatum and Sesarma angolense as burrowers and Armases elegans as a climber. The non-burrowing Perisesarma alberti and the potentially phytothelmic Sesarma buettikoferi (Fusi et al. unpublished data, Fig. 4B), on the other hand, were more dense in the Avicennia belt of Wouri Bridge forest (with an average of about 4 individuals per square meter) the former and in the *Pandanus* belt in the same forest (almost 6 individuals per square meter) for the latter. Within molluscs, we recorded the presence of the Thiarid Pachymelania fusca in the two forests with a density of at about 400 specimens per square meter and the Potamidid Tympanotonus radula (Fig. 4A) found only in Bois de Singes forest with a density of at about 40 individuals per square meter.

A significant difference in macrobenthos assemblages between Wouri Bridge and Bois des Singes forest was recorded (F = 25.655, p = 0.01, Figs. 3 and 4). Specifically, Bois des Singe was characterized by the absence of *S. buettikoferi* and *S. angolense* and a more evenly distributed species density. In Wouri Bridge forest, a dominant species for each belt was observed: *P. alberti* was dominant in the *Avicennia* belt, *S. buettikoferi* in the *Pandanus* belt, while *P. alberti* and *P. kamermanni* were



Fig. 2. Average concentrations of Sterols (A), PAHs (B) and metals (C), detected in sediment samples. Data are shown according to vegetation belt (*Avicennia* sp., *Pandanus* sp., *Rhizophora* sp.) and sampling depth on the x-axis. Surface sediment upper layer (0–10 cm) and core samples from 11 to 20 cm.

the two most abundant species in the *Rhizophora* belt. There was a notable absence of the gastropod *T. radula* throughout all the Wouri Bridge transects.

#### Table 1

Total PCB concentrations in sediment samples. Sample size is shown in brackets. Data are expressed as mean  $\pm$  standard error.

Forest	Belt	Depth (cm)	PCB tot (µg kg^{-1})
Bois des singes	Rhizophora	10 (n = 7)	$6.2\pm1.9$
		20 (n = 7)	$6.3 \pm 1.8$
Wouri Bridge	Avicennia	10(n = 3)	$6.3 \pm 0.4$
		20(n = 3)	$5.0 \pm 3$
	Pandanus	10(n = 3)	$2.6 \pm 1.2$
		20(n = 3)	-
	Rhizophora	10 (n = 3)	$5.3 \pm 1.5$
		20(n = 3)	$4.0 \pm 1.2$

# 3.3. Macrobenthos assemblages and contaminants

The DistLM analysis shows a significant relationship between macrobenthos assemblage and sterols, metals, PAHs and C/N data (Table 5; Fig. 5A) and explains >90% of the total variation. In particular, 5  $\beta$ -cholestan-3 $\alpha$ -ol l, Selenium, Chromium and Zinc explain the highest percentage of variation (20, 15, 8 and 7% respectively, p < 0.01). The two species of mollusc, the crabs *Perisesarma huzardi* and *P. alberti* appear to be most affected by the variation of the significant environmental variables cited above (Fig. 5B).

# 4. Discussion

Mangroves in Cameroon still cover many hectares of estuaries, especially along the Wouri River where a complex system of channels and

# 728 **Table 2**

p-p'DDE and DEHP concentrations in sediment samples. Sample size is shown in brackets. Data are expressed as mean  $\pm$  standard error.

Forest	Belt	Depth (cm)	p,p'- DDE ( $\mu g \ kg^{-1}$ )	DEHP ( $\mu g \ kg^{-1}$ )
Bois des singes	Rhizophora	10 (n = 7)	$30\pm0.2$	$750\pm110$
		20 (n = 7)	$10 \pm 0.1$	$540 \pm 90$
Wouri Bridge	Avicennia	10(n = 3)	-	$710\pm370$
		20 (n = 3)	-	$530 \pm 110$
	Pandanus	10 (n = 3)	-	$380 \pm 40$
		20 (n = 3)	-	$1008\pm390$
	Rhizophora	10 (n = 3)	$40\pm0.07$	$750\pm290$
		20 (n = 3)	$40\pm0.07$	$970\pm60$

fens hinder access, and thus direct exploitation, mainly represent by logging for household and coal, and land claim for building new settlements. Nevertheless, the rapid development of Douala together with important commercial and trade activities, due to the presence of the harbour, have contributed to the city being a source of contaminants, which are spreading into nearby mangrove forests. We revealed in this study the presence of contaminants such as PCBs, PAHs, DEHP and heavy metals.

# 4.1. Effect of sterols, p,p'-DDE and heavy metals contamination on macrobenthic fauna

In particular, p-p-DDE (DDT metabolite) was found in high concentrations close to the threshold admissible limit (Yang et al., 2007), as

previously reported in other mangrove systems by Bodin et al. (2011) and Bhupander and Debapriya (2012). It is likely that this high level is related to recent intensive, and mainly uncontrolled, anti-malaria treatment in the area (Antonio-Nkondjio et al., 2011; Denison, 2013; Etang et al., 2007; Fossog Tene et al., 2013). Unfortunately, these compounds are reported to have a toxic effect on marine organisms (e.g. Bayarri et al., 2001; Mearns et al., 2014) and we strongly suggest that their high levels found in Wouri Bridge might be the reason for the total lack of Tympanotonus radula (Fig. 4, Table 2) since the documented endocrine disrupting mechanism of p,p'-DDE in molluscs (Matthiessen, 2008). Extremely rapid urbanization has resulted in a growing urban population that has colonised areas within well-established rainforest and mangrove forests. In these areas, settlements consist of rudimentary housing with uncontrolled discharge of untreated sewage and wastewater into the forests (Nfotabong-Atheull et al., 2011; Simon and Raffaelli, 2012). This is likely the cause of the presence of sterols detected with a high ratio of (Coprostanol + epicoprostanol)/ $\sum$  Total Stanols that indicates a serious level of sediment sewage contamination (Froehner et al., 2009; Gern and da C. Lana, 2013). Specifically, we identified high contamination in 10 of the 26 samples (4 in Wouri Bridge and 6 in Bois des Singes). This contamination is determined not only by the uncontrolled urbanization taking place in Douala, but also by the lack of any wastewater treatment management in the city. Indeed, the area nearby Bois de Singes is highly affected by municipality wastewater discharge, with trucks (personal observation) releasing tons of untreated wastewater directly into the mangrove, also witnessed during the survey. This activity is also a good explanation of the fact that we found more

#### Table 3

Stanol contamination index: percentage of coprostan-3-ol and 5  $\beta$ -cholestan-3 $\alpha$ -ol on total sterols, calculated for the different depths in each belt of the two forests. High levels of contaminants (> 25% according with Froehner et al., 2009) are shown in bold.

Samples	Site	Depth	Belt	% (coprostan-3-ol $+5\beta\text{-cholestan-3}\alpha\text{-ol}$ )/total stanols
A1PW10	Wouri Bridge	10	Avicennia	43.9
A1PW20	Wouri Bridge	20	Avicennia	0.1
P1PW10	Wouri Bridge	10	Pandanus	9.8
P1PW20	Wouri Bridge	20	Pandanus	0.3
P2PW10	Wouri Bridge	10	Pandanus	47.8
P2PW20	Wouri Bridge	20	Pandanus	5.4
R1PW10	Wouri Bridge	10	Rhizophora	6.3
R1PW20	Wouri Bridge	20	Rhizophora	7.1
R2PW10	Wouri Bridge	10	Rhizophora	44.5
R2PW20	Wouri Bridge	20	Rhizophora	3.3
R3PW10	Wouri Bridge	10	Rhizophora	67.4
R3PW20	Wouri Bridge	20	Rhizophora	9.0
R1BS10	Bois des singes	10	Rhizophora	24.0
R1BS20	Bois des singes	20	Rhizophora	3.7
R2BS10	Bois des singes	10	Rhizophora	11.9
R2BS20	Bois des singes	20	Rhizophora	21.6
R3BS10	Bois des singes	10	Rhizophora	43.1
R3BS20	Bois des singes	20	Rhizophora	39.1
R4BS10	Bois des singes	10	Rhizophora	23.0
R4BS20	Bois des singes	20	Rhizophora	25.6
R5BS10	Bois des singes	10	Rhizophora	21.5
R5BS20	Bois des singes	20	Rhizophora	35.6
R6BS10	Bois des singes	10	Rhizophora	26.0
R6BS20	Bois des singes	20	Rhizophora	14.1
R7BS10	Bois des singes	10	Rhizophora	68.2
R7BS20	Bois des singes	20	Rhizophora	18.8

Ratio values that represent detection of contamination are shown in bold. BS = Bois des Singes; PW = Wouri Bridge.

#### Table 4

Sediment water temperature, pH AND Conductivity and pore water total nitrogen (TN) and organic carbon (TOC) recorded in each plot during macrobenthos surveys. Data are expressed as mean  $\pm$  standard error.

Forest	Belt	Temperature (°C)	рН	Conductivity (mV)	Tot N (%)	Tot OC (%)
Bois des singes	Rhizophora	$27.35\pm0.3$	$6.29 \pm 0.06$	$16.13\pm2.9$	$0.24\pm0.04$	$4.09\pm0.49$
Wouri Bridge	Avicennia	$26.28 \pm 0.4$	$6.25\pm0.04$	$20.83 \pm 2.3$	$0.28\pm0.09$	$4.55\pm1.38$
	Pandanus	$26.9 \pm 0.4$	$6.30\pm0.03$	$18.08 \pm 1.6$	$0.33 \pm 0.2$	$5.65\pm0.19$
	Rhizophora	$26.18\pm0.06$	$6.29 \pm 0.07$	$19 \pm 2$	$0.38\pm0.1$	$6.01 \pm 1.25$



**Fig. 3.** Non-metric multidimensional scaling ordination showing the patterns of distribution of macrobenthic species in the two study forests. ♦ Bois des Singes *Rhizophora*; ● Wouri Bridge *Avicennia*; + Wouri Bridge *Pandanus*; ♦ Wouri Bridge *Rhizophora*.

highly contaminated samples in Bois de Singes than Wouri Bridge. Through the macrobenthic survey, we were able to record highly biodiverse and structured macrobenthic communities in both forests. However, we recorded two significantly different patterns of macrobenthic assemblage at the two sites, mainly due to the absence of *T. radula* (Potamididae) and *P. huzardi* (Sesarmidae) in Wouri Bridge forest, while *S. buettikoferi* was not found in Bois de Singes forest (Fig. 4). Our statistical analyses found 5  $\beta$ -cholestan-3 $\alpha$ -ol (Gern and da C. Lana, 2013) and four heavy metals, As (Beltman et al., 1999), Se (Hamilton, 2004), Cr (Lewis et al., 2001) and Zn (Ellis et al., 2004; Schaffelke et al., 2005), to be the main drivers of differences in crab assemblage (See Fig. 5 and Table 5). Indeed, they are known to be among the more important compounds responsible to disrupt the physiology of marine species (e.g. Mello and Nayak, in press and reference above). If we consider that the main environmental features, including the tidal regime, were similar between sites and belts, we hypothesize that these contaminant levels could play a major role shaping the macrobenthos distribution and density as a result of their differential sensitivity to pollutants and their concentration. Such a selective ecological effect has been largely described for molluscs in east African mangrove systems exposed to sewage (Cannicci et al., 2009; Penha-Lopes et al., 2010).

## 4.2. Implications of anthropogenic contamination on mangrove ecosystem

Currently, the interpretation of biological responses as a consequence of contamination remains complex. A major reason being that organisms in the field are exposed to multiple stressors under dynamic conditions (e.g. variable micro- and macronutrient loads, changing climatic conditions, multiple contaminants, tidal cycles and salinity), and potential additive, synergistic or antagonistic responses to these stressors may occur (Bayen, 2012). Indeed, changes in the diversity/ structure of mangrove ecosystems have been reported as a response to chemical pollution (e.g. Mohamed et al., 2008), which has also been linked to a decline in some populations such as mangrove oysters and snails (e.g. Roach and Wilson, 2009) and molluscs (Cannicci et al., 2009). Kulkarni et al. (2010) reported low biodiversity indices associated with a low water quality index in mangrove ecosystems in India. The overall impact of pollution, however, appears to be complex. For



Fig. 4. Densities of mollusc (A) and crab (B) species in the study sites. Values are expressed as mean  $\pm$  SE.

#### Table 5

Test for relationships between sterols, metals, PAHs and C/N with macrobenthos distribution, using permutational multiple regression analysis (DISTLM). AICc: coefficient of regression, SS: sum of squares, F: value of pseudo and its significance p (% Var: percentage of variance explained by each single variable, and % Cumul: cumulative percentage of variance explained, Res. df: residual degrees of freedom). In bold the variable statistically significant related with macrobenthos distribution.

Variable	AICc	SS(trace)	Pseudo-F	р	% Var.	%Cumul.	Res.df
+ coprostan-3-ol	0.012	113.38	0.29	0.7352	0.001	0.012	24
+ 5 $\beta$ -cholestan-3 $\alpha$ -ol	0.219	1978.7	6.12	0.0105	0.208	0.219	23
+ cholesterol	0.291	678.86	2.21	0.1209	0.071	0.291	22
+ 5 $\alpha$ -cholestan-3 $\beta$ -ol	0.314	219.05	0.70	0.4644	0.023	0.314	21
+ As	0.434	1151.3	4.27	0.0316	0.121	0.434	20
+ Se	0.587	1458.1	7.05	0.0073	0.153	0.587	19
+ Mo	0.604	155.92	0.74	0.471	0.016	0.604	18
+ Cd	0.625	197.66	0.94	0.3786	0.021	0.625	17
+ Sn	0.637	119.91	0.55	0.5553	0.013	0.637	16
+ Sb	0.669	300.56	1.43	0.241	0.032	0.669	15
+ Cr	0.751	785.68	4.64	0.0322	0.082	0.751	14
+ Cu	0.804	508.87	3.55	0.0613	0.053	0.804	13
+ Mn	0.809	45.498	0.30	0.7313	0.005	0.809	12
+ Zn	0.885	717.9	7.18	0.0095	0.075	0.885	11
+ phenanthrene	0.900	149.18	1.57	0.2309	0.016	0.900	10
+ fluoranthene	0.901	6.8011	0.06	0.8834	0.001	0.901	9
+ benzo[a] anthracene	0.918	166.41	1.71	0.2141	0.017	0.918	8
+ benzo [jbk]fluoranthene	0.922	38.665	0.37	0.71	0.004	0.922	7
+ benzo [a] pyrene	0.922	-3.598	-0.03	0.9607	0.000	0.922	6
+ DEHP	0.968	85.979	1.13	0.3724	0.009	0.968	4
+ p-p'DDE	0.984	149.9	2.91	0.1425	0.016	0.984	3
+ C/N	0.990	56.641	1.16	0.365	0.006	0.990	2

example, the patterns of diversity and species composition recorded in various mangrove forests highly impacted by humans in Indonesia did not clearly correlate with the impact investigated (Geist et al., 2012).

Moreover, discharge of domestic sewage at low levels caused an increase in crab population size in east Africa (Cannicci et al., 2009; Penha-Lopes et al., 2011) and did not affect the macrobenthic



**Fig. 5.** Distance-based redundancy analysis plots (dbRDA) of macrobenthos distribution across Wouri Bridge (*Avicennia* belt ( $\bullet$ ), *Pandanus* belt (+) and *Rhizophora* belt ( $\bullet$ )) and Bois des Singes ( $\diamond$ ), in accordance with the contaminants found in the sediment core of each belt. Vectors correspond to environmental variables (A) and species (B). Length and direction of the vectors indicate the strength of the correlation between the variable and ordination axis given the other variables in the model. The radius of the circle denotes a correlation of 1.

communities in Hong Kong (Wong et al., 1997; Yu et al., 1997). Our results indicate that multiple anthropogenic stressors, and in particular heavy loads of wastewater, although not resulting in the hypothesised depletion of crab abundance in Wouri River estuary mangroves, can shape their community composition. The data suggest that contaminant loadings in Wouri river mangroves, affecting the distribution of macrobenthonic species, could lead to the type of cryptic ecological degradation (sensu Dahdouh-Guebas et al. 2005) shown by Bartolini et al. (2011), who documented an inverse relationship between the increased biomass of fiddler crabs and their overall engineering function, thus affecting the whole mangrove ecosystem.

The sterols level recorded in this study strongly indicates a more large contamination by sewage and, in particular, by a first stages of human faecal contamination, as recorded in other studies (i.e. Fernandes et al. 1999; Silva & Madureira 2012). This can have an important effect on the ecology of macrobenthic species (Frena et al., 2016) by changing the chemistry of the sediment where they live dramatically enhancing bacterial activity (Dheenan et al., 2016). The subtle faecal contamination could triggering possible ecological bottom-up effect with relevant ecological consequences on the overall system (Hamilton, 2004), through the modification of secondary consumers primary feeding sources, such as microbenthic (Isobe et al., 2004) or infaunal macrobenthic communities (Moon et al., 2008). Together with the presence of PCBs, PAHs and heavy metals, chromium in particular, raises at least two important concerns. The first is that mangrove sediment receiving wastewater does accumulate a remarkably higher level of long-term contaminants (Tam and Wong, 1995). The second is that they could potentially spread through the whole mangrove area (Agoramoorthy et al., 2008; Yi et al., 2011; Zhou et al., 2007), ultimately affecting, through the trophic chain, the secondary and tertiary consumers which are consumed by the local population. Therefore, they potentially represent a health risk.

#### 5. Conclusions

To our knowledge, this study is the first to investigate the contamination of the Cameroon estuarine and marine environment with Persistent Organic Pollutants (POPS). It contributes to the scarce data and literature on this subject in African countries, which has mainly focused on public health and resistance implications from POPs use. Our data clearly show that the main source of contamination in the mangrove forests surrounding Douala is represented by uncontrolled discharge of urban wastewater and the persistent, illegal and indiscriminate use of DDT. These contaminants, together with four specific heavy metals (As, Cr, Zn, Se) seem to affect the macrobenthonic assemblage of the two study sites, suggesting that Douala peri-urban mangrove is subjected to a complex patchwork of contamination. This documented inflow pollution has serious implications for ecosystem functioning and public health. Therefore we emphasize the necessity to prioritise water quality monitoring and the development of public policies for the wastewater management. Additionally, as highlighted in many studies, human pollution is likely to impair the provision of critical mangrove ecosystem services which are relied upon by local communities. Integrated assessment of macrobenthic assemblages should be considered as a method to detect early contamination patterns, as suggested by our results and confirmed by several other studies. Hence, the present data provide a baseline for further development and environmental management oriented towards anthropogenic pollution by POPs in west Africa, also in the view of monitoring and reducing human impact to mitigate vulnerability of mangroves to the fast climate change.

# Acknowledgements

The study was supported by the Cameroon Oil Transportation Company (COTCO), the SP3-People (Marie Curie) IRSES Project CREC (no. 247514), the SC research funds from MIUR (ex 60%) and supported by King Abdullah University of Science and Technology (baseline research funds to DD). We thank Adolphe Nfotabong-Atheull, Vanessa Ngo-Massou and Joseph Bayi for invaluable help during field sampling. We are grateful to Ilaria Marchi for design assistance and Pierluisa Fantini for invaluable laboratory work. Thanks to Jenny Marie Booth for English revision of this manuscript.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at http://dx. doi.org/10.1016/j.marpolbul.2016.06.104.

#### References

- Agoramoorthy, G., Chen, F., Hsu, M.J., 2008. Threat of heavy metal pollution in halophytic and mangrove plants of Tamil Nadu, India. Environ. Pollut. 155, 320–326. http://dx. doi.org/10.1016/j.envpol.2007.11.011.
- Alemagi, D., 2007. The oil industry along the Atlantic coast of Cameroon: assessing impacts and possible solutions. Resour. Policy 32, 135–145. http://dx.doi.org/10.1016/ j.resourpol.2007.08.007.
- Anderson, M.J., Godley, R.N., Clarke, K.R., 2008. PERMANOVA + for PRIMER: Guide to Software and Statistical Methods. PRIMER-E Ltd., Plymouth.
- Andreetta, A., Fusi, M., Cameldi, I., Cimò, F., Carnicelli, S., Cannicci, S., 2014. Mangrove carbon sink. Do burrowing crabs contribute to sediment carbon storage? Evidence from a Kenyan mangrove system. J. Sea Res. 85, 524–533. http://dx.doi.org/10.1016/j. seares.2013.08.010.
- Antonio-Nkondjio, C., Fossog, B.T., Ndo, C., Djantio, B.M., Togouet, S.Z., Awono-Ambene, P., Costantini, C., Wondji, C.S., Ranson, H., 2011. Anopheles gambiae distribution and insecticide resistance in the cities of Douala and Yaoundé (Cameroon): influence of urban agriculture and pollution. Malar. J. 10, 154. http://dx.doi.org/10.1186/1475-2875-10-154.
- Ball, M.C., 1988. Ecophysiology of mangroves. Trees 2, 129–142. http://dx.doi.org/10. 1007/BF00196018.
- Bartolini, F., Cimò, F., Fusi, M., Dahdouh-Guebas, F., Lopes, G.P., Cannicci, S., 2011. The effect of sewage discharge on the ecosystem engineering activities of two East African fiddler crab species: consequences for mangrove ecosystem functioning. Mar. Environ. Res. 71, 53–61. http://dx.doi.org/10.1016/j.marenvres.2010.10.002.
- Bayarri, S., Baldassarri, L.T., Iacovella, N., Ferrara, F., Domenico, A.D., 2001. PCDDs, PCDFs, PCBs and DDE in edible marine species from the Adriatic Sea. Chemosphere 43, 601–610. http://dx.doi.org/10.1016/S0045-6535(00)00412-4.
- Bayen, S., 2012. Occurrence, bioavailability and toxic effects of trace metals and organic contaminants in mangrove ecosystems: a review. Environ. Int. 48, 84–101. http:// dx.doi.org/10.1016/j.envint.2012.07.008.
- Bayen, S., Wurl, O., Karuppiah, S., Sivasothi, N., Lee, H.K., Obbard, J.P., 2005. Persistent organic pollutants in mangrove food webs in Singapore. Chemosphere 61, 303–313. http://dx.doi.org/10.1016/j.chemosphere.2005.02.097.
- Bayen, S., Estrada, E., Juhel, G., Lee, W., Kelly, B., 2016. Pharmaceutically active compounds and endocrine disrupting chemicals in water, sediments and mollusks in mangrove ecosystems from Singapore. Mar. Pollut. Bull. 109, 716–722.
- Beltman, D.J., Clements, W.H., Lipton, J., Cacela, D., 1999. Benthic invertebrate metals exposure, accumulation, and community-level effects downstream from a hard-rock mine site. Environ. Toxicol. Chem. 18, 299–307. http://dx.doi.org/10.1897/1551-5028(1999)018<0299:bimeaa>2.3.co;2.
- Berti, R., Cannicci, S., Fabbroni, S., Innocenti, G., 2008. Notes on the structure and the use of Neosarmatium meinerti and Cardisoma Carnifex burrows in a Kenyan mangrove swamp (Decapoda Brachyura). Ethol. Ecol. Evol. 20, 101–113.
- Bettinelli, M., Beone, G., Spezia, S., Baffi, C., 2000. Determination of heavy metals in soils and sediments by microwave-assisted digestion and inductively coupled plasma optical emission spectrometry analysis. Anal. Chim. Acta 424, 289–296. http://dx.doi. org/10.1016/S0003-2670(00)01123-5.
- Bhupander, K., Debapriya, M., 2012. Eco-toxicological risk assessment of HCH, DDT and their possible sources by isomeric ratio distribution in sediments from sundarban mangrove ecosystem in Bay of Bengal, India. J. Environ. Earth Sci. 2, 58–71.
- Bodin, N., N'Gom Ka, R., Le Loc'h, F., Raffray, J., Budzinski, H., Peluhet, L., Tito de Morais, L., 2011. Are exploited mangrove molluscs exposed to persistent organic pollutant contamination in Senegal, West Africa? Chemosphere 84, 318–327. http://dx.doi.org/10. 1016/j.chemosphere.2011.04.012.
- Bull, I.D., Lockheart, M.J., Elhmmali, M.M., Roberts, D.J., Evershed, R.P., 2002. The origin of faeces by means of biomarker detection. Environ. Int. 27, 647–654. http://dx.doi.org/ 10.1016/S0160-4120(01)00124-6.
- Campo, E.V., Darius, M., 2004. Mangrove palynology in recent marine sediments off Cameroon. Mar. Geol. 208, 315–330. http://dx.doi.org/10.1016/j.margeo.2004.04.014.
- Cannicci, S., Burrows, D., Fratini, S., Smith, T.J., Offenberg, J., Dahdouh-Guebas, F., 2008. Faunal impact on vegetation structure and ecosystem function in mangrove forests: a review. Aquat. Bot. 89, 186–200. http://dx.doi.org/10.1016/j.aquabot.2008.01.009.
- Cannicci, S., Bartolini, F., Dahdouh-Guebas, F., Fratini, S., Litulo, C., Macia, A., Mrabu, E.J., Penha-Lopes, G., Paula, J., 2009. Effects of urban wastewater on crab and mollusc assemblages in equatorial and subtropical mangroves of East Africa. Estuar. Coast. Shelf Sci. 84, 305–317. http://dx.doi.org/10.1016/j.ecss.2009.04.021.
- Dafforn, K.a., Simpson, S.L., Kelaher, B.P., Clark, G.F., Komyakova, V., Wong, C.K.C., Johnston, E.L., 2012. The challenge of choosing environmental indicators of

anthropogenic impacts in estuaries. Environ. Pollut. 163, 207–217. http://dx.doi.org/10.1016/j.envpol.2011.12.029.

- Dahdouh-Guebas, F., Hettiarachchi, S., Lo Seen, D., Batelaan, O., Sooriyarachchi, S., Jayatissa, L.P., Koedam, N., 2005. Transitions in ancient inland freshwater resource management in Sri Lanka affect biota and human populations in and around coastal lagoons. Curr. Biol. 15, 579–586. http://dx.doi.org/10.1016/j.cub.2005.01.053.
- Denison, L, 2013. Stockholm Convention on persistent organic pollutants. Air Qual. Clim. Chang. 2.
- Dheenan, P.S., Jha, D.K., Das, A.K., Vinithkumar, N.V., Devi, M.P., Kirubagaran, R., 2016. Geographic information systems and multivariate analysis to evaluate fecal bacterial pollution in coastal waters of Andaman, India. Environ. Pollut. 214, 45–53. http://dx. doi.org/10.1016/j.envpol.2016.03.065.
- Din, N., Baltzer, F., 2008. Richesse Floristique et Evolution des mangroves de l'Estuaire du Cameroun. Afr. Geosci. Rev. 2, 119–130.
- Din, N., Priso, R.J., Kenne, M., Ngolio, D.E., Blasco, F., 2002. Early growth stages and natural regeneration of *Avicennia germinans* (L.) stearn in the Wouri estuarine mangroves (Douala-Cameroon). Wetl. Ecol. Manag. 10, 461–472. http://dx.doi.org/10.1023/A: 1021351707822.
- Diop, S., Barusseau, J., Descamps, C., 2014. The land/ocean interactions in the coastal zone of West and Central Africa, Springer, estuaries of the world. Springer International Publishing, Cham http://dx.doi.org/10.1007/978-3-319-06388-1.
- Duke, N.C., 2016. Oil spill impacts on mangroves: a global review with recommendations. Mar. Pollut. Bull. 109, 700–715.
- Duke, N., Meynecke, J.O., Dittmann, S., Ellison, A.M., Anger, K., Berger, U., Cannicci, S., Diele, K., Ewel, K.C., Field, C.D., Koedam, N., Lee, S.Y., Marchand, C., Nordhaus, I., Dahdouh-Guebas, F., 2007. A world without mangroves ? Science 317, 41–43. http://dx.doi. org/10.1126/science.317.5834.41b (80-. ).
- Ellis, J., Nicholls, P., Craggs, R., Hofstra, D., Hewitt, J., 2004. Effects of terrigenous sedimentation on mangrove physiology and associated macrobenthic communities. Mar. Ecol. Prog. Ser. 270, 71–82. http://dx.doi.org/10.3354/meps270071.
- Ellison, J.C., Zouh, I., 2012. Vulnerability to climate change of mangroves: assessment from Cameroon, Central Africa. Biology (Basel) 1, 617–638. http://dx.doi.org/10.3390/ biology1030617.
- Emmerson, W.D., 2001. Aspects of the population dynamics of *Neosarmatium meinerti* at Mgazana, a warm temperate mangrove swamp in the East Cape, South Africa, investigated using an indirect method 221–229.
- Etang, J., Manga, L., Toto, J., Guillet, P., Fondjo, E., Etang, J., Manga, L., Toto, J., Guillet, P., Fondjo, E., 2007. Spectrum of metabolic-based resistance to DDT and pyrethroids in *Anopheles gambiae* sl. populations from Cameroon spectrum of metabolic-based resistance to DDT and pyrethroids in *Anopheles gambiae* sl populations from Cameroon. Vector Ecol. 32, 123–133.
- Fattore, E., Benfenati, E., Marelli, R., Cools, E., Fanelli, R., 1996. Sterols in sediment samples from Venice Lagoon, Italy. Chemosphere 33, 2383–2393. http://dx.doi.org/10.1016/ S0045-6535(96)00340-2.
- Fernandes, M.B., Sicre, M.a., Cardoso, J.N., Macêdo, S.J., 1999. Sedimentary 4-desmethyl sterols and n-alkanols in an eutrophic urban estuary, Capibaribe River, Brazil. Sci. Total Environ. 231, 1–16. http://dx.doi.org/10.1016/S0048-9697(99)00077-7.
- Fossog Tene, B., Poupardin, R., Costantini, C., Awono-Ambene, P., Wondji, C.S., Ranson, H., Antonio-Nkondjio, C., 2013. Resistance to DDT in an urban setting: common mechanisms implicated in both M and S forms of *Anopheles gambiae* in the city of Yaound?? Cameroon. PLoS One 8. http://dx.doi.org/10.1371/ journal.pone.0061408.
- Fratini, S., Cannicci, S., Vannini, M., 2000. Competition and interaction between Neosarmatium smithi (Crustacea : Grapsidae) and Terebralia palustris (Mollusca : Gastropoda) in a Kenyan mangrove. Mar. Biol. 137, 309–316.
- Frena, M., Bataglion, G.A., Tonietto, A.E., Eberlin, M.N., Alexandre, M.R., Madureira, L.A.S., 2016. Assessment of anthropogenic contamination with sterol markers in surface sediments of a tropical estuary (Itajaí-Açu, Brazil). Sci. Total Environ. 544, 432–438. http://dx.doi.org/10.1016/j.scitotenv.2015.11.137.
- Friess, D.A., Webb, E.L., 2014. Variability in mangrove change estimates and implications for the assessment of ecosystem service provision. Glob. Ecol. Biogeogr. 23, 715–725. http://dx.doi.org/10.1111/geb.12140.
- Froehner, S., Martins, R.F., Errera, M.R., 2009. Assessment of fecal sterols in Barigui River sediments in Curitiba, Brazil. Environ. Monit. Assess. 157, 591–600. http://dx.doi. org/10.1007/s10661-008-0559-0.
- Gabche, C.E., 1997. An Appraisal of Fisheries Activities and Evaluation of Economic Potential of the Fish Trade in the Douala–Edea Reserve–Cameroon, Cameroon W. ed. Yaoundé, Cameroon.
- Geist, S.J., Nordhaus, I., Hinrichs, S., 2012. Occurrence of species-rich crab fauna in a human-impacted mangrove forest questions the application of community analysis as an environmental assessment tool. Estuar. Coast. Shelf Sci. 96, 69–80. http://dx. doi.org/10.1016/j.ecss.2011.10.002.
- Gern, F.R., da C. Lana, P., 2013. Reciprocal experimental transplantations to assess effects of organic enrichment on the recolonization of benthic macrofauna in a subtropical estuary. Mar. Pollut. Bull. 67, 107–120. http://dx.doi.org/10.1016/j.marpolbul.2012. 11.026.
- Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J., Duke, N., 2011. Status and distribution of mangrove forests of the world using earth observation satellite data. Glob. Ecol. Biogeogr. 20, 154–159. http://dx.doi.org/10.1111/j.1466-8238. 2010.00584.x.
- Hamilton, S.J., 2004. Review of selenium toxicity in the aquatic food chain. Sci. Total Environ. 326, 1–31. http://dx.doi.org/10.1016/j.scitotenv.2004.01.019.
- Isobe, K.O., Tarao, M., Chiem, N.H., Minh, L.Y., 2004. Effect of environmental factors on the relationship between concentrations of coprostanol and Fecal Indicator bacteria in tropical (Mekong Delta) and temperate (Tokyo) freshwaters. Appl. Environ. Microbiol. 70, 814–821. http://dx.doi.org/10.1128/AEM.70.2.814.

- Kristensen, E., 2008. Mangrove crabs as ecosystem engineers; with emphasis on sediment processes. J. Sea Res. 59, 30–43. http://dx.doi.org/10.1016/j.seares.2007. 05.004.
- Kulkarni, V.A., Jagtap, T.G., Mhalsekar, N.M., Naik, A.N., 2010. Biological and environmental characteristics of mangrove habitats from Manori creek, West Coast, India. Environ. Monit. Assess. 168, 587–596. http://dx.doi.org/10.1007/s10661-009-1136-x.
- Lee, S.Y., 2008. Mangrove macrobenthos: assemblages, services, and linkages. J. Sea Res. 59, 16–29. http://dx.doi.org/10.1016/j.seares.2007.05.002.
- Lewis, M.A., Weber, D.E., Stanley, R.S., Moore, J.C., 2001. Dredging impact on an urbanized Florida bayou: effects on benthos and algal-periphyton. Environ. Pollut. 115, 161–171. http://dx.doi.org/10.1016/S0269-7491(01)00118-X.
- Lewis, M., Pryor, R., Wilking, L. 2011. Fate and effects of anthropogenic chemicals in mangrove ecosystems: a review. Environ. Pollut. 159, 2328–2346. http://dx.doi.org/10. 1016/j.envpol.2011.04.027.
- Lovelock, C.E., Cahoon, D.R., Friess, D.A., Guntenspergen, G.R., Krauss, K.W., Reef, R., Rogers, K., Saunders, M.L., Sidik, F., Swales, A., Saintilan, N., Thuyen, L.X., Triet, T., 2015. The vulnerability of Indo-Pacific mangrove forests to sea-level rise. Nature 526, 559-U217. http://dx.doi.org/10.1038/nature15538.
- MacFarlane, G.R., Koller, C.E., Blomberg, S.P., 2007. Accumulation and partitioning of heavy metals in mangroves: a synthesis of field-based studies. Chemosphere 69, 1454–1464. http://dx.doi.org/10.1016/j.chemosphere.2007.04.059.
- Manning, R.B., Holthuis, L.B., 1981. West African Brachyuran crabs (Crustacea: Decapoda). Smithson. Contrib. Zool. 1–396 http://dx.doi.org/10.5479/si.00810282.306.
- Massó i Alemán, S., Bourgeois, C., Appeltans, W., Vanhoorne, B., De Hauwere, N., Stoffelen, P., Heughebaert, A., Dahdouh-Guebas, F., 2010. The "mangrove reference database and herbarium.". Plant Ecol. Evol. 143, 225–232. http://dx.doi.org/10.5091/plecevo. 2010.439.
- Matthiessen, P., 2008. An assessment of endocrine disruption in mollusks and the potential for developing internationally standardized mollusk life cycle test guidelines. Integr. Environ. Assess. Manag. 4, 274–284. http://dx.doi.org/10.1897/IEAM\_2008-003.1.
- Mearns, A.J., Reish, D.J., Oshida, P.S., Ginn, T., Rempel-Hester, M.A., Arthur, C., Rutherford, N., 2014. Effects of pollution on marine organisms. Water Environ. Res. 86, 1869–1954. http://dx.doi.org/10.2175/106143012X13407275695751.
- Mello, C.A.N., Nayak, G.N., 2016. Assessment of metal enrichment and their bioavailability in sediment and bioaccumulation by mangrove plant pneumatophores in a tropical ( Zuari) estuary, west coast of India. Mar. Pollut. Bull. http://dx.doi.org/10.1016/j. marpolbul.2016.06.0590025-326X (in press).
- Mohamed, M.O.S., Neukermans, G., Kairo, J.G., Dahdouh-Guebas, F., Koedam, N., 2008. Mangrove forests in a peri-urban setting: the case of Mombasa (Kenya). Wetl. Ecol. Manag. 17, 243–255. http://dx.doi.org/10.1007/s11273-008-9104-8.
- Moon, H.B., Yoon, S.P., Jung, R.H., Choi, M., 2008. Wastewater treatment plants (WWTPs) as a source of sediment contamination by toxic organic pollutants and fecal sterols in a semi-enclosed bay in Korea. Chemosphere 73, 880–889. http://dx.doi.org/10.1016/j. chemosphere.2008.07.038.
- Mudge, S.M., Bebianno, M.J.a.F., East, J.a., Barreira, L.a., 1999. Sterols in the Ria Formosa lagoon, Portugal. Water Res. 33, 1038–1048. http://dx.doi.org/10.1016/S0043-1354(98)00283-8.
- Nfotabong-Atheull, A., Din, N., Longonje, S.N., Koedam, N., Dahdouh-Guebas, F., 2009. Commercial activities and subsistence utilization of mangrove forests around the Wouri estuary and the Douala-Edea reserve (Cameroon). J. Ethnobiol. Ethnomed. 5, 35. http://dx.doi.org/10.1186/1746-4269-5-35.
- Nfotabong-Atheull, A., Din, N., Essomè Koum, LG., Satyanarayana, B., Koedam, N., Dahdouh-Guebas, F., 2011. Assessing forest products usage and local residents' perception of environmental changes in peri-urban and rural mangroves of Cameroon, Central Africa. J. Ethnobiol. Ethnomed. 7, 41. http://dx.doi.org/10.1186/1746-4269-7-41.
- Nfotabong-Atheull, A., Din, N., Dahdouh-Guebas, F., 2013. Qualitative and quantitative characterization of mangrove vegetation structure and dynamics in a peri-urban setting of Douala (Cameroon): an approach using air-borne imagery. Estuar. Coasts 36, 1181–1192. http://dx.doi.org/10.1007/s12237-013-9638-8.
- Ng, P.K.L., Davie, P.J.F., Guinot, D., 2008. Systema Brachyurorum : part I. An annotated checklist of extant Brachyuran crabs of the world. Raffles Bull. Zool. 17, 1–286.
- Ngo-Massou, V., Essome-Koum, G., Ngollo-Dina, E., Din, N., 2012. Composition of macrobenthos in the Wouri River estuary mangrove, Douala, Cameroon. Afr. J. Mar. Sci. 34, 349–360. http://dx.doi.org/10.2989/1814232X.2012.725288.
- Peng, X., Zhang, G., Mai, B., Hu, J., Li, K., Wang, Z., 2005. Tracing anthropogenic contamination in the Pearl River estuarine and marine environment of South China Sea using sterols and other organic molecular markers. Mar. Pollut. Bull. 50, 856–865. http:// dx.doi.org/10.1016/j.marpolbul.2005.02.031.
- Penha-Lopes, G., Kristensen, E., Flindt, M., Mangion, P., Bouillon, S., Paula, J., 2010. The role of biogenic structures on the biogeochemical functioning of mangrove constructed wetlands sediments – a mesocosm approach. Mar. Pollut. Bull. 60, 560–572. http:// dx.doi.org/10.1016/j.marpolbul.2009.11.008.
- Penha-Lopes, G., Torres, P., Cannicci, S., Narciso, L., Paula, J., 2011. Monitoring anthropogenic sewage pollution on mangrove creeks in southern Mozambique: a test of *Palaemon concinnus* Dana, 1852 (Palaemonidae) as a biological indicator. Environ. Pollut. 159, 636–645. http://dx.doi.org/10.1016/j.envpol.2010.09.029.
- Peters, E.C., Gassman, N.J., Firman, J.C., Richmond, R.H., Power, E.a., 1997. Ecotoxicology of tropical marine ecosystems. Environ. Toxicol. Chem. 16, 12–40. http://dx.doi.org/10. 1002/etc.5620160103.
- Price, A., Klaus, A., Sheppard, C.R.C., Abbiss, M.A., Kofani, M., Webster, G., 2000. Environmental and bioeconomic characterisation of coastal and marine systems of Cameroon, including risk implications of the Chad–Cameroon pipeline project. Aquat. Ecosyst. Health Manag. 3, 137–161. http://dx.doi.org/10.1016/S1463-4988(99)00072-X.

- Roach, A.C., Wilson, S.P., 2009. Ecological impacts of tributyltin on estuarine communities in the Hastings River, NSW Australia. Mar. Pollut. Bull. 58, 1780–1786. http://dx.doi. org/10.1016/j.marpolbul.2009.08.021.
- Saenger, P., Bellan, M.F., 1995. The Mangrove Vegetation of the Atlantic Coast of Africa. Universite' de Toulouse Press, Toulouse.
- Schaffelke, B., Mellors, J., Duke, N.C., 2005. Water quality in the Great Barrier Reef region: responses of mangrove, seagrass and macroalgal communities. Mar. Pollut. Bull. 51, 279–296. http://dx.doi.org/10.1016/j.marpolbul.2004.10.025.
- Sherwin, M.R., Van Vleet, E.S., Fossato, V.U., Dolci, F., 1993. Coprostanol (5β-cholestan-3βol) in lagoonal sediments and mussels of Venice, Italy. Mar. Pollut. Bull. 26, 501–507. http://dx.doi.org/10.1016/0025-326X(93)90467-X.
- Silva, C.a., Madureira, L.a.S., 2012. Source correlation of biomarkers in a mangrove ecosystem on Santa Catarina Island in southern Brazil. An. Acad. Bras. Cienc. 84, 589–604.
- Simon, L.N., Raffaelli, D., 2012. Assessing ecosystem effects of small-scale cutting of Cameroon mangrove forests. J. Ecol. Nat. Environ. 4, 126–134. http://dx.doi.org/10.5897/ JENE11.131.
- Skov, M.W., Vannini, M., Shunula, J.P., Hartnoll, R.G., Cannicci, S., 2002. Quantifying the density of mangrove crabs: Ocypodidae and Grapsidae. Mar. Biol. 141, 725–732.
- Sousa, W.P., Dangremond, E.M., 2011. Trophic interactions in coastal and estuarine mangrove forest ecosystems. Treatise on Estuarine and Coastal Science. Academic London, Waltham, pp. 43–93.
- Spalding, M., Kainuma, M., Collins, L., 2010. World Atlas of Mangroves. Earthscan, p. 319. Sparks, D.L., Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T., Sumner, M.E., 1996. Methods of soil analysis. Part 3 - Chemical Methods. Soil Science Society of America Inc., Madison, WI.
- Tam, N.F.Y., Wong, Y.S., 1995. Spatial and temporal variations of heavy metal contamination in sediments of a mangrove swamp in Hong Kong. Mar. Pollut. Bull. 31, 254–261. http://dx.doi.org/10.1016/0025-326X(95)00141-9.
- UNEP, 2007. Mangroves of Western and Central Africa. Earthprint, Chicago.

- Van De Walle, N., 1989. rice Politics in Cameroon: state commitment, capability, and urban bias. J. Mod. Afr. Stud. 27, 579. http://dx.doi.org/10.1017/S0022278X00020450.
- Vane, C.H., Harrison, I., Kim, A.W., Moss-Hayes, V., Vickers, B.P., Hong, K., 2009. Organic and metal contamination in surface mangrove sediments of South China. Mar. Pollut. Bull. 58, 134–144. http://dx.doi.org/10.1016/j.marpolbul.2008.09.024.
- Wong, S., Tam, N.F.Y., Lan, C.Y., 1997. Mangrove wetlands as wastewater treatment facility : a field trial. Hydrobiologia 352, 49–59.
- Woodroffe, C.D., Rogers, K., McKee, K.L., Lovelock, C.E., Mendelssohn, I.a., Saintilan, N., 2016. Mangrove sedimentation and response to relative sea-level rise. Ann. Rev. Mar. Sci. 8. http://dx.doi.org/10.1146/annurev-marine-122414-034025 (annurevmarine-122414-034025).
- Yang, J., Zhang, W., Shen, Y., Feng, W., Wang, X., 2007. Monitoring of organochlorine pesticides using PFU systems in Yunnan lakes and rivers, China. Chemosphere 66, 219–225. http://dx.doi.org/10.1016/j.chemosphere.2006.05.055.
- Yi, Y., Yang, Z., Zhang, S., 2011. Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze River basin. Environ. Pollut. 159, 2575–2585. http://dx.doi. org/10.1016/j.envpol.2011.06.011.
- Yu, R., Chen, G.Z., Wong, Y.S., Tam, N.F.Y., Lan, C.Y., 1997. Benthic macrofauna of the mangrove swamp treated with municipal wastewater. Hydrobiologia 347, 127–137.
- Zaccone, C., Gallipoli, A., Cocozza, C., Trevisan, M., Miano, T.M., 2009. Distribution patterns of selected PAHs in bulk peat and corresponding humic acids from a Swiss ombrotrophic bog profile. Plant Soil 315, 35–45. http://dx.doi.org/10.1007/s11104-008-9775-1.
- Zhou, F., Guo, H., Hao, Z., 2007. Spatial distribution of heavy metals in Hong Kong's marine sediments and their human impacts: a GIS-based chemometric approach. Mar. Pollut. Bull. 54, 1372–1384. http://dx.doi.org/10.1016/j.marpolbul.2007.05.017.